



A New Diploid Number and Sex Chromosome System for the Family Hersiliidae from the Mediterranean Region of Turkey

Ümit Kumbıçak, Zübeyde Kumbıçak* and Gülnare Huseynli

¹Department of Molecular Biology and Genetics, Science and Art Faculty, Nevşehir Hacı Bektaş Veli University, Nevşehir, Turkey

ABSTRACT

Cytogenetic features, mitotic metaphase chromosomes, karyogram and course of meiosis of *Hersiliola bayrami* were determined based on Giemsa-stained testicular chromosomes for the first time. Results of spermatogonial metaphases showed that the chromosome number and the karyotype formula of the species was $2n♂=35 (X_1X_2X_30)$ which is the first record for the family Hersiliidae. Both autosomes and sex chromosomes were acrocentric. The lengths of the chromosomes indicated no significant difference between autosomal pairs, however, X_3 was the smallest chromosome but X_1 and X_2 were intermediate in the karyotype.

Article Information

Received 03 March 2017

Revised 25 May 2017

Accepted 03 July 2017

Available online 02 January 2018

Authors' Contribution

UK collected spiders, helped in laboratory work, made karyotype and analysed the data. ZK helped in laboratory, field trips and wrote and evaluated the manuscript. GH made chromosome slides and arranged the measurements of chromosomes.

Key words

Araneae, Cytogenetics, *Hersiliola*, Karyotype, Spider.

INTRODUCTION

Hersiliidae Thorell, 1870 is a small family, currently including 181 species belonging to 16 genera (World Spider Catalog, 2017). The hersiliids are conspicuously long-legged, delicate spiders, characterised by extremely elongated posterior lateral spinnerets (Baehr and Baehr, 1987). The majority of hersiliid species are found in tropical and subtropical regions. Recently, they are a subject of several revisions of Australian, Oriental, Neotropical, and Afrotropical faunas. In the Palaearctic, *Hersiliola* Thorell, 1870 is the most species rich genus (Marusik *et al.*, 2010).

Cytogenetical data on spiders are important from the perspective of understanding their evolution. Karyological information such as chromosome number and the morphology and size of chromosomes is useful in deducing evolution and interrelationships in the group (Joshi *et al.*, 2016). Diploid chromosome numbers of spiders range from 7 (Suzuki, 1954) to 128 (Král *et al.*, 2013). Entelegynae spiders (the most derived group) exhibit lower diploid numbers and mostly monoarmed chromosomes when compared with the predominantly high chromosome numbers and biarmed chromosomes of mygalomorphs (Kořínková and Král, 2013; Kumbıçak *et al.*, 2014). A multiple sex chromosome system of the X_1X_20 type is the most common in spiders (*i.e.*, Suzuki, 1954) but is rare

in other animal groups. In many spider species, sex is not determined by presence of an Y or W chromosome, as it occurs in most mammals and birds (Araújo *et al.*, 2016). Compared with this system, $X_1X_2X_30$ sex chromosome system can rarely be seen in some spider families (*i.e.*, Agelenidae; Ctenidae; Eresidae; Eutichuridae; Linyphiidae; Nemesiidae; Oecobiidae; Selenopidae; Sparassidae; Tetragnathidae) (Araújo *et al.*, 2017).

In the present study, we aimed to analyze the chromosomes of an endemic species, *Hersiliola bayrami* Danişman *et al.*, 2012 for which chromosomal data were lacking.

MATERIALS AND METHODS


A sample of nine adult males of *H. bayrami* were analyzed. They were collected from the Mediterranean part of Turkey during March-May in 2014 (Table I). The specimens were deposited in the collection of Genetics Laboratory, Science and Art Faculty, Molecular Biology and Genetics Department, Nevşehir.

The chromosomal preparations were made according to the method of Pekár and Král (2001). The gonads were dissected out in a physiological solution for invertebrate animals and incubated for 12-15 min in hypotonic solution (0.0075 M KCl); immediately after the gonads were fixed in a freshly prepared methanol (or absolute ethanol): acetic acid (3:1) for 10 and 20 min. The fixed tissues were put in a drop of 60% acetic acid and macerated using a tungsten needle on a histological plate (42°C).

* Corresponding author: zkumbicak@nevsehir.edu.tr
0030-9923/2018/0001-0131 \$ 9.00/0
Copyright 2018 Zoological Society of Pakistan

Table I.- Number of used specimens, collection data including locality, geographical coordinates and dates.

Species	Sample	Collection data	
		Locality (districts) and geographical coordinates	Date
<i>Hersiliola bayrami</i> Danışman <i>et al.</i> , 2012	2♂♂	Alanya, Antalya; 36°32'16.84" N and 31°59'33.90"E	21.04.2014
	1♂	Gözne, Mersin; 37°00'30.47" N and 34°31'50.09"E	28.03.2014
	2♂♂	Anamur, Mersin, 36°05'59.38" N and 32°49'47.17"E	02.05.2014
	3♂♂	Anamur, Mersin, 36°02'17.48" N and 32°44'00.54"E	04.05.2014
	1♂	Centre, Karaman, 37°08'45.07" N and 33°13'30.92"E	17.04.2014

Fig. 1. Karyotype of *Hersiliola bayrami* based on male specimens. Scale bar=10 μ m.Fig. 2. *Hersiliola bayrami* ($2n♂=35, X_1X_2X_3$). A, Spermatogonial metaphase; B, Pachytene; C, Late diakinesis; D, Late metaphase I; E, Metaphase II; F, Anaphase II (arrows sex chromosomes; asterisks, nuclei with sex chromosomes). Scale bars=10 μ m.

Slides were investigated under a phase contrast microscope (Leica DM2500) and the slides containing well figures were further analyzed. The slides were stained with 5% Giemsa solution in Sørensen's phosphate buffer (pH=6.8) for 27 min. Mitotic and meiotic stages were photographed by a BX53 light microscope (Olympus) equipped with DP26 digital camera with CellSens software (Olympus). Karyotypes were constructed by arranging chromosomes in pairs according to size using images of spermatogonial metaphases. Relative chromosome lengths (RCLs) of each chromosome pair were calculated from 10 metaphases. Chromosome morphology was classified according to the nomenclature proposed by Levan *et al.* (1964).

RESULTS

The male karyotype of *H. bayrami* consisted of $2n♂=35$ chromosomes (Fig. 2A). All chromosome pairs were acrocentric including the sex chromosomes. The sex chromosome system was $♂X_1X_2X_3/♀X_1X_1X_2X_2X_3X_3$. Values for the RCLs were between 7.94% to 4.23% and gradually decreased in size. RCL of X_1 and X_2 was 6.63% and 5.02%, respectively. The X_3 was the smallest element in the karyotype (Fig. 1).

At the stages of meiotic prophase I (*i.e.*, leptotene, zygotene and pachytene), the sex chromosomes were positively heteropycnotic and large X_1 and X_2 associated with each other. However, the small X_3 chromosome was located close to X_1 and X_2 but did not associate with them (Fig. 2B). During diplotene, diakinesis and metaphase I, 16 autosomal bivalents and three univalent sex chromosomes were seen; generally, one chiasma (*i.e.*, interstitial or terminal type) per bivalent was observed (Fig. 2C, D). During the second meiotic stages (*i.e.*, prophase II, metaphase II and anaphase II), sex chromosomes were indistinguishable because they were isopycnotic with the autosomes. Two types of nuclei were observed, either with ($n=19$) or without ($n=16$) sex chromosomes (Fig. 2E, F).

DISCUSSION

A karyotype of only one hersiliid species (*Hersilia savignyi* Lucas, 1836) was described up to now. Most of authors reported the diploid number as $2n♂=30$ for the species (Bole-Gowda, 1958; Srivastava and Shukla, 1986; Parida and Sharma, 1987; Sharma and Parida, 1987; Prakash and Prakash, 2014), however, Sharma *et al.* (1960) reported also $2n♂=32$. The sex chromosome system was always of the $♂X_1X_2/♀X_1X_1X_2X_2$ type. Karyotype of the genus *Hersiliola* was reported for the first time by this study with the male diploid number $2n♂=35$ (32,

$X_1X_2X_3$) which is not so common in spiders. Hersiliidae is closely related with the family Oecobiidae, where the known diploid numbers listed in Araújo *et al.* (2017) are $2n♂=22$ in *Oecobius cellariorum* Dugès, 1836 (Youju *et al.*, 1993), $2n♂=25$ in *O. putus* O. Pickard-Cambridge, 1876 (Mittal, 1983; Srivastava and Shukla, 1986), $2n♂=39$ in *Uroctea lesserti* Schenkel, 1936 (Youju *et al.*, 1993), $2n♂=42$ in *U. compactilis* L. Koch, 1878 (Suzuki, 1950, 1954). Based on these results; there is no strong relation on diploid numbers between families, nevertheless, the sex chromosome type $X_1X_2X_3$ is encountered in both families Hersiliidae and Oecobiidae. To explain the chromosomal differences between these families, additional cytogenetic studies are needed including the species not investigated previously.

The chromosomes in the family studied to date were acrocentric (Araújo *et al.*, 2017), however, Prakash and Prakash (2014) reported also metacentric, submetacentric and subtelocentric chromosomes in the karyotype of *H. savignyi*. In our study, all autosomes and sex chromosomes of *H. bayrami* were acrocentric. This may be due to differences in the techniques used to obtain the figures of chromosomes and measurements of them.

Our study reported the first results of the chromosome number, karyotype and meiotic features for the genus *Hersiliola*. The species in focus, *H. bayrami* was investigated by a classical cytogenetic analysis. This study will promote further studies of the family Hersiliidae.

ACKNOWLEDGEMENTS

We are thankful to Dr. Osman Seyyar (Department of Biology, Art and Science Faculty, Ömer Halisdemir University, Niğde, Turkey) for the first spider determination and Serhat Bayrak (B.S., Department of Biology, Art and Science Faculty, Nevşehir Hacı Bektaş Veli University, Nevşehir, Turkey) for breeding the spiders.

Statement of conflict of interest

Authors have declared no conflict of interest.

REFERENCES

- Araújo, D., Schneider, M.C., Paula-Neto, E. and Cella, D.M., 2017. *The spider cytogenetic database, Version 5.5*. Available at: www.arthropodacytogenetics.bio.br/spiderdatabase (accessed 17 Jan 2017).
- Araújo, D., Sanches, M.B., Lima, J.S.G.S., Nascimento, E.V.J., Giroti, A.M., Brescovit, A.D., Cella, D.M. and Schneider, M.C., 2016. Chromosomal analyses of Salticinae and Lyssomaninae reveal a broad

- occurrence of the $2n\sigma = 28, X_1X_2,0$ karyotype within Salticidae. *J. Arachnol.*, **44**: 148-152. <https://doi.org/10.1636/M15-77>
- Baehr, B. and Baehr, M., 1987. The Australian Hersiliidae (Arachnida Araneae): Taxonomy, Phylogeny, Zoogeography. *Inverteb. Taxon*, **1**: 351-437. <https://doi.org/10.1071/IT9870351>
- Bole-Gowda, B.N., 1958. A study of the chromosomes during meiosis in twenty-two species of Indian spiders. *Proc. zool. Soc. Bengal*, **11**: 69-108.
- Joshi, H.S., Yadav, P.B., Lekhak, M.M. and Yadav, S.R., 2016. Cytogenetics of two endemic *Barleria* species (Acanthaceae) from the northern Western Ghats (India). *Caryologia*, **69**: 170-174. <https://doi.org/10.1080/00087114.2016.1152111>
- Kořínková, T. and Král, J., 2013. Karyotypes, sex chromosomes and meiotic division in spiders. In: *Spider ecophysiology* (ed. W. Netwig). Springer-Verlag, Berlin, pp. 159-171. https://doi.org/10.1007/978-3-642-33989-9_12
- Král, J., Kořínková, T., Krkavcová, L., Musilová, J., Forman, M., Herrera, I.M.A., Haddad, C.R., Vítková, M., Henriques, S., Vargas, J.G.P. and Hedin, M., 2013. Evolution of karyotype, sex chromosomes, and meiosis in mygalomorph spiders (Araneae: Mygalomorphae). *Biol. J. Linn. Soc.*, **109**: 377-408. <https://doi.org/10.1111/bij.12056>
- Kumbıçak, Z., Ekiz, E. and Çiçekli, S., 2014. Karyotypes of six spider species belonging to the families Gnaphosidae, Salticidae, Thomisidae, and Zodariidae (Araneae) from Turkey. *Comp. Cytogene.*, **8**: 93-101. <https://doi.org/10.3897/compcytogen.v8i2.6065>
- Levan, A., Fredga, K. and Sandberg, A.A., 1964. Nomenclature for centromeric position on chromosomes. *Hereditas*, **52**: 201-220.
- Marusik, Y.M., Kunt, K.B. and Yağmur, E.A., 2010. A new species of *Hersiliola* Thorell, 1870 (Araneae, Hersiliidae) from Turkey. *ZooKeys*, **37**: 27-34. <https://doi.org/10.1111/j.1601-5223.1964.tb01953.x>
- Mittal, O.P., 1983. Karyological studies on the Indian spiders X. An XXX0-type of sex-determining mechanism in the cribellate spiders. *La Kromosomo II*, **30-31**: 919-925.
- Parida, B.B. and Sharma, N.N., 1987. Chromosome number, sex mechanism and genome size in 27 species of Indian spiders. *Chromos. Inform. Serv.*, **43**: 11-13.
- Pekár, S. and Král, J., 2001. A comparative study of the biology and karyotypes of two central European Zodariid spiders (Araneae, Zodariidae). *J. Arachnol.*, **29**: 345-353. [https://doi.org/10.1636/0161-8202\(2001\)029\[0345:ACSOTB\]2.0.CO;2](https://doi.org/10.1636/0161-8202(2001)029[0345:ACSOTB]2.0.CO;2)
- Prakash, A. and Prakash, S., 2014. Cytogenetical investigations on spiders of semi-arid areas. *Ind. J. of Arachnol.*, **3**: 40-54.
- Sharma, N. and Parida, B.B., 1987. Study of chromosomes in spiders from Orissa. *Pranikee*, **8**: 71-76.
- Sharma, G.P., Tandon, K.K. and Grewal, M.S., 1960. Cytological studies on the Indian spiders V. chromosome complement and male meiosis in *Hersilia savignyi* Lucas (Hersiliidae), *Larinia* sp. (Argiopidae), *Tetragnatha* sp. (Tetragnathidae), *Oxyopes ryvesii* Pocock, and *Oxyopes* sp. (Oxyopidae). *Res. Bull. Panjab Univ.*, **11**: 201-206.
- Srivastava, M.D.L. and Shukla, S., 1986. Chromosome number and sex-determining mechanism in forty-seven species of Indian spiders. *Chromos. Inform. Serv.*, **41**: 23-26.
- Suzuki, S., 1950. Spiders with extremely low and high chromosome numbers. *Jpn. J. Genet.*, **25**: 221-222.
- Suzuki, S., 1954. Cytological studies in spiders. III. Studies on the chromosomes of fifty-seven species of spiders belonging to seventeen families, with general considerations on chromosomal evolution. *J. Sci. Hiroshima Univ. B*, **15**: 23-136.
- World Spider Catalog, 2017. *The World spider catalog*, Version 18.0. Natural History Museum Bern. Available at: <http://wsc.nmbe.ch> (accessed 7 Jun 2017).
- Youju, W., Daxiang, S., Xiuzhen, W. and Zhenling, Y., 1993. Preliminary studies on the chromosome of four species of spiders. *Acta Arachnol. Sin.*, **2**: 110-113.